

CH7 EXAM - REVIEW QUESTION - Solutions

$$f(x) = x^3/4 - x^2/3 - x/2 + 3 \cos x$$

EQUATION OF line L : $L(x) = f(a) + f'(a)(x-a)$

$$f(0) = 3$$

$$f'(x) = \frac{3}{4}x^2 - \frac{2}{3}x - \frac{1}{2} - 3 \sin x$$

$$f'(0) = -\frac{1}{2}$$

$$L(x) = 3 - \frac{1}{2}x$$

a) $f(x) = 0$ @ $x = -1.373, x = 1.346, x = 3.164$

$$A_R = \int_{-1.373}^0 f(x) dx = 2.903$$

b) $V = \pi \int_{-1.373}^0 [(f(x)+2)^2 - (0+2)^2] dx \approx 59.361$

c) $A_{\Delta} = \frac{1}{2}(\text{base})\left(\frac{\sqrt{3}}{2}\text{base}\right) = \frac{\sqrt{3}}{4}(\text{base})^2$

$$V = \int_{-1.373}^0 \frac{\sqrt{3}}{4} (f(x))^2 dx \quad \text{or} \quad \frac{\sqrt{3}}{4} \int_{-1.373}^0 (f(x))^2 dx$$

$$\rightarrow \int(3)/4 * \int((y1(x))^{\wedge}2, x, -1.373, 0)$$

CALC SYNTAX, IF asked to solve.

d) $P = \underbrace{1.373}_{\text{along } x} + \underbrace{3}_{\text{along } y} + \int_{-1.373}^0 \sqrt{1+[f'(x)]^2} dx$ * $f'(x) = \frac{3}{4}x^2 - \frac{2}{3}x - \frac{1}{2} - 3 \sin x$

3.390 ← intersection

* NOTE: you must state what $f'(x)$ equals somewhere on the paper. ~~None~~ is written in


e) $A_s = \int_0^{\underbrace{3-\frac{1}{2}x}_{L(x)}} [L(x) - f(x)] dx$

f) Key: $g'(x) = f(x)$ [FTC-1]

$$L = \int_{-1}^0 \sqrt{1+[g'(x)]^2} dx = \int_{-1}^0 \sqrt{1+[f(x)]^2} dx \approx 2.792$$

Calc Syntax * $\int(1+(y1(x))^{\wedge}2), x, -1, 0)$

→ TAKE A WHILE TO COMPUTE !! ←



2 (NO CALCULATOR) Consider the curves $y = e^x$, $y = x^3$, $y = 5 \ln(x-1)$, and $y = 4$ each shown above.

Let R1 be the light shaded region bounded in the first quadrant by the graphs of $y = e^x$, $y = x^3$, and $y = 4$.

Let R2 be the dark shaded region bounded in the first quadrant by the graphs of $y = x^3$, $y = 5 \ln(x-1)$, and $y = 4$.

A. Write, but do not evaluate, an expression involving one or more integrals used to find the volume of the solid generated when R2 is rotated about the vertical line $x = 6$.

B. The region R2 is the base of a solid. Every cross section perpendicular to the y -axis is a square whose side lies flat on R2. Write, but do not evaluate, an expression involving one or more integrals that can be used to find the volume of the solid.

C. **FIND** the area of region R2.

D. Write, but do not evaluate, an expression involving one or more integrals, used to find the area of region R1.

E. Write, but do not evaluate, an expression involving one or more integrals, used to find the perimeter of region R2.

Solutions on next page

2 $y = e^x \rightarrow x = \ln y$ $y = x^3 \rightarrow x = \sqrt[3]{y}$ $y = 5 \ln(x-1) \rightarrow x = e^{y/5} + 1$

A.

Washers: $V = \pi \int_0^4 [(6 - \sqrt[3]{y})^2 - (6 - (e^{y/5} + 1))^2] dy$

(OR) Cylindrical Shells: $V = \int_0^{\sqrt[3]{4}} 2\pi(6-x)[x] dx + \int_{\sqrt[3]{4}}^2 2\pi(6-x)[4] dx + \int_2^{1+e^{4/5}} 2\pi(6-x)[4] dx$

B. $A_{\square} = (\text{base})^2 = (e^{y/5} + 1 - \sqrt[3]{y})^2$

$V = \int_0^4 A dy = \int_0^4 \underbrace{[e^{y/5} + 1 - \sqrt[3]{y}]^2}_{\text{base}} dy$

C. $\int_0^4 [e^{y/5} + 1 - \sqrt[3]{y}] dy = \left[5e^{y/5} + y - \frac{3}{4}y^{4/3} \right]_0^4 = (5e^{4/5} + 4 - \frac{3}{4}(4)^{4/3}) - (5)$

D. $\Delta x: A = \int_0^{\ln 4} [e^x - x^3] dx + \int_{\ln 4}^{\sqrt[3]{4}} [4 - x^3] dx$

(OR) $\Delta y: A = \int_0^1 \sqrt[3]{y} dy + \int_1^4 [\sqrt[3]{y} - \ln y] dy$

E. $\Delta x: P = \underbrace{2}_{\text{Side 1}} + \underbrace{\int_2^{1+e^{4/5}} \sqrt{1 + (\frac{5}{x-1})^2} dx}_{\text{Side 2}} + \underbrace{[(e^{4/5} + 1) - \sqrt[3]{4}]}_{\text{Side 3}} + \underbrace{\int_0^{\sqrt[3]{4}} \sqrt{1 + (3x^2)^2} dx}_{\text{Side 4}}$

(OR)

$\Delta y: P = \underbrace{2}_{\text{Side 1}} + \underbrace{\int_0^4 \sqrt{1 + (\frac{1}{5}e^{y/5})^2} dy}_{\text{Side 2}} + \underbrace{e^{4/5} + 1 - \sqrt[3]{4}}_{\text{Side 3}} + \underbrace{\int_0^4 \sqrt{1 + (\frac{1}{3y^{2/3}})^2} dy}_{\text{Side 4}}$